




Introduction to Machine Learning

Machine Learning: Jordan Boyd-Graber
University of Maryland


SLACK SVMS

Slides adapted from Eric Xing

Can SVMs Work Here?




Can SVMs Work Here?



$$y_i(w \cdot x_i + b) \geq 1 \quad (1)$$

Trick: Allow for a few bad apples



New objective function

$$\min_{w, b, \xi} \frac{1}{2} \|w\|^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

New objective function

$$\min_{w, b, \xi} \frac{1}{2} \|w\|^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

- Standard margin

New objective function

$$\min_{w, b, \xi} \frac{1}{2} \|w\|^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

- Standard margin
- How wrong a point is (slack variables)

New objective function

$$\min_{w, b, \xi} \frac{1}{2} \|w\|^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

- Standard margin
- How wrong a point is (slack variables)
- Tradeoff between margin and slack variables

New objective function

$$\min_{w, b, \xi} \frac{1}{2} \|w\|^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$


- Standard margin
- How wrong a point is (slack variables)
- Tradeoff between margin and slack variables
- **How bad wrongness scales**

Aside: Loss Functions

- Losses measure how bad a mistake is
- Important for slack as well


Aside: Loss Functions

- Losses measure how bad a mistake is
- Important for slack as well




Aside: Loss Functions

- Losses measure how bad a mistake is
- Important for slack as well




Aside: Loss Functions

- Losses measure how bad a mistake is
- Important for slack as well



Aside: Loss Functions

- Losses measure how bad a mistake is
- Important for slack as well



We'll focus on linear hinge loss

Optimizing Constrained Functions

Theorem: Lagrange Multiplier Method

Given functions $f(x_1, \dots, x_n)$ and $g(x_1, \dots, x_n)$, the critical points of f restricted to the set $g = 0$ are solutions to equations:

$$\begin{aligned}\frac{\partial f}{\partial x_i}(x_1, \dots, x_n) &= \lambda \frac{\partial g}{\partial x_i}(x_1, \dots, x_n) \quad \forall i \\ g(x_1, \dots, x_n) &= 0\end{aligned}$$

This is $n + 1$ equations in the $n + 1$ variables x_1, \dots, x_n, λ .

Lagrange Example

Maximize $f(x, y) = \sqrt{xy}$ subject to the constraint $20x + 10y = 200$.

- Compute derivatives

Lagrange Example

Maximize $f(x, y) = \sqrt{xy}$ subject to the constraint $20x + 10y = 200$.

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2} \sqrt{\frac{y}{x}}$$

Lagrange Example

Maximize $f(x, y) = \sqrt{xy}$ subject to the constraint $20x + 10y = 200$.

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2} \sqrt{\frac{y}{x}} \quad \frac{\partial g}{\partial x} = 20$$

Lagrange Example

Maximize $f(x, y) = \sqrt{xy}$ subject to the constraint $20x + 10y = 200$.

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2} \sqrt{\frac{y}{x}} \quad \frac{\partial g}{\partial x} = 20$$

$$\frac{\partial f}{\partial y} = \frac{1}{2} \sqrt{\frac{x}{y}}$$

Lagrange Example

Maximize $f(x, y) = \sqrt{xy}$ subject to the constraint $20x + 10y = 200$.

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2} \sqrt{\frac{y}{x}} \quad \frac{\partial g}{\partial x} = 20$$

$$\frac{\partial f}{\partial y} = \frac{1}{2} \sqrt{\frac{x}{y}} \quad \frac{\partial g}{\partial y} = 10$$

Lagrange Example

Maximize $f(x, y) = \sqrt{xy}$ subject to the constraint $20x + 10y = 200$.

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2} \sqrt{\frac{y}{x}} \quad \frac{\partial g}{\partial x} = 20$$

$$\frac{\partial f}{\partial y} = \frac{1}{2} \sqrt{\frac{x}{y}} \quad \frac{\partial g}{\partial y} = 10$$

- Create new systems of equations

$$\frac{1}{2} \sqrt{\frac{y}{x}} = 20\lambda$$

$$\frac{1}{2} \sqrt{\frac{x}{y}} = 10\lambda$$

$$20x + 10y = 200$$

Lagrange Example

- Dividing the first equation by the second gives us

$$\frac{y}{x} = 2 \quad (3)$$

- which means $y = 2x$, plugging this into the constraint equation gives:

$$20x + 10(2x) = 200$$

$$x = 5 \Rightarrow y = 10$$

New Lagrangian

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i(\mathbf{w} \cdot \mathbf{x}_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

New Lagrangian

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i(\mathbf{w} \cdot \mathbf{x}_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients ($\nabla_{\mathbf{w}} \mathcal{L}, \nabla_b \mathcal{L}, \nabla_{\xi_i} \mathcal{L}$) and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7)$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i \quad (8)$$

$$\alpha_i + \beta_i = C \quad (9)$$

New Lagrangian

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i(\mathbf{w} \cdot \mathbf{x}_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients ($\nabla_{\mathbf{w}} \mathcal{L}$, $\nabla_b \mathcal{L}$, $\nabla_{\xi_i} \mathcal{L}$) and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7)$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i \quad (8)$$

$$\alpha_i + \beta_i = C \quad (9)$$

New Lagrangian

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i(\mathbf{w} \cdot \mathbf{x}_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients ($\nabla_{\mathbf{w}} \mathcal{L}$, $\nabla_b \mathcal{L}$, $\nabla_{\xi_i} \mathcal{L}$) and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7)$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i \quad (8)$$

$$\alpha_i + \beta_i = C \quad (9)$$

New Lagrangian

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i(\mathbf{w} \cdot \mathbf{x}_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients ($\nabla_{\mathbf{w}} \mathcal{L}$, $\nabla_b \mathcal{L}$, $\nabla_{\xi_i} \mathcal{L}$) and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7)$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i \quad (8)$$

$$\alpha_i + \beta_i = C \quad (9)$$

Simplifying dual objective

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i x_i$$

$$\alpha_i + \beta_i = C$$

Simplifying dual objective

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i x_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \|\vec{w}\|^2 - \sum_i \alpha_i y_i \vec{w} \cdot \vec{x}_i - \sum_i \alpha_i y_i b - \sum_{i=1}^m \beta_i \xi_i \quad (10)$$

Simplifying dual objective

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b - \sum_{i=1}^m \beta_i \xi_i \quad (10)$$

Simplifying dual objective

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b - \sum_{i=1}^m \beta_i \xi_i \quad (10)$$

Simplifying dual objective

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b + \sum_i^m \alpha_i \quad (10)$$

Simplifying dual objective

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b + \sum_i^m \alpha_i \quad (10)$$

Simplifying dual objective

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b + \sum_i^m \alpha_i \quad (10)$$

Simplifying dual objective

$$\begin{aligned} \sum_{i=1}^m \alpha_i y_i &= 0 & \vec{w} &= \sum_{i=1}^m \alpha_i y_i x_i & \alpha_i + \beta_i &= C \\ \mathcal{L} &= \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) + \sum_i^m \alpha_i \end{aligned} \quad (10)$$

First two terms are the same!

Simplifying dual objective

$$\begin{aligned} \sum_{i=1}^m \alpha_i y_i &= 0 & \vec{w} &= \sum_{i=1}^m \alpha_i y_i x_i & \alpha_i + \beta_i &= C \\ \mathcal{L} &= -\frac{1}{2} \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) + \sum_i^m \alpha_i & & & & (10) \end{aligned}$$

Just like separable case, except that we add the constraint that $\alpha_i \leq C$!

Wrapup

- Adding slack variables don't break the SVM problem
- Very popular algorithm
 - SVMLight (many options)
 - Libsvm / Liblinear (very fast)
 - Weka (friendly)
 - pyml (Python focused)